

Crestal sinus lift using a turbine fitted with spherical diamond-faced burs with stops: Non-traumatic technique

MA Lopez¹, M Andreasi-Bassi¹, L Confalone¹, F Carinci^{2*}

Abstract

Introduction

One of the most frequent reasons for failures during the operation of maxillary sinus floor lift is connected with the possibility of a rupture to the Schneiderian membrane which, if lacerated, cannot perform its function of graft containment. In order to reduce the incidence of complications the turbine is frequently used in association with spherical diamond-faced burs to cut the hard tissue with extreme accuracy and minor trauma, while saving the soft tissue.

In this study a new technique of maxillary crestal sinus lift performed with the aid of a turbine with spherical diamond-faced burs is proposed. Calibrated stops have been created to perfectly control the procedure.

Case Study

The patient was a female, 47 years old, non-smoker, healthy with no systemic disease, not on regular prescribed medication, periodontally treated, who needed an operation of maxillary sinus lift. The cortical of the maxillary sinus is reduced through the use of these burs so as to obtain a hole which can enable both access to the maxillary sinus and, subsequently, the lifting of the Schneiderian membrane. Working time is reduced to less than 3 minutes in the cortical thinning operation and percussive trauma is avoided.

Conclusion

Due to the reduction in trauma and invasiveness of the process, this technique could be a valid alternative

to the techniques known and applied to date.

Introduction

With the increasing demand for implant-prosthetic treatments, there is a similar increase in the requirement for anatomic and morpho-structural sites for the application of fixtures. The anatomic limitations reduce the possibility to perform standard treatments and consequently the operator needs advanced surgical techniques in order to resolve more complex cases.

Often in the upper distal edentulous maxilla a marked osseous re-absorption is found, with large pneumatization of the maxillary sinus. This condition requires surgical lifting techniques of the maxillary sinus floor by means of which it is possible to transform part of the sinus cavity into osseous tissue adequate for implant-prosthesis.

Lifting the maxillary sinus floor can be achieved through different techniques thanks to osseous inlays: part of the maxillary sinus is filled with biocompatible material which after some months are transformed into bone thanks to re-absorption and remodelling processes.

When large cavity fillings are required, it is generally necessary to resort to vestibular approaches to the maxillary sinus. In these cases it is not always possible to insert the implants at the same time as the lifting procedure. A second operation is required when there is a residual osseous thickness, between three and six mm, such as to allow a good primary stability for the fixtures.

The crestal approach is increasingly employed where small quantities of

biomaterial are required to fill the sinus cavity. In these cases it is generally possible to insert the implants immediately. These 'mini lifts' are therefore advisable when the initial osseous thickness is between 5 and 6 mm¹.

The techniques employed for the mini lift of the maxillary sinus are various, and for more than thirty years they have been continuously improving.

Among the first employed techniques was the Favero-Branemark technique that foresaw a minimum of 7-8 mm initial osseous height, utilising a greenwood (bone distraction) fracture of the sinus cortical. Despite the success of this technique only modest increments of bone growth, up to a maximum of 1.5-2 mm, were achieved².

One among the most widespread techniques is the Summers compaction technique, which is primarily used in the sites where there is low-density bone (D3, D4).

This technique prescribes the employment of compacting osteotomes with increasing diameter^{3,4,5}.

The main advantage is the resulting bone compaction, frequently not very thick in the upper distal sectors, while the disadvantage is due to the numerous and traumatic percussions. The lift can reach 3 - 4 mm.

The same technique was later modified in 1999-2002 by Fugazzotto, who introduced the employment of a trephine drill^{6,7,8,9,10}.

In 2000 Cosci¹¹ tried to make it even less traumatic by using drills without sharp (working) points.

*Corresponding author
Email: crc@unife.it

¹ Private professional in Rome

² Chair of Maxillofacial Surgery, Department of Morphology, Surgery and Experimental Medicine, University of Ferrara

A variation of Summers expansion technique is the alternate osteotomic technique proposed by Malchiodi, where first concave and then convex chisels are used alternatively with increasing diameters¹².

A very widespread method is the one proposed by Zaninari that requires the sinus floor fracture with blunted chisels. The fragment of the fractured cortical is then used to lift the sinus membrane and subsequently the area is filled with biomaterial. Such a technique, that does not foresee compaction, reduces but does not eliminate the percussive trauma in comparison with Summers' technique.

While it reduces the number of chisels, the percussion strength remains a risk for the integrity of the Schneiderian membrane¹³.

Recently there is the tendency to mainly develop techniques with crestal approaches such as the trans-alveolar lift with or without graft material and contemporaneous crest expansion that seem, in some cases, to achieve similar results to those obtained with vestibular access¹⁴.

The main reason for failure of these operations is connected to the membrane laceration with consequent dispersion of the graft material in the maxillary sinus and therefore a failure in osseous reconstitution around the implant apex.

Another negative element is represented by the traumatism and invasiveness of these operations that often require various percussive procedures, resulting in trauma. An interesting proposal is given by the ultrasound approach that slightly reduces trauma^{15,16,17,18,19,20}. In particular, ultrasounds crestal approach seems to be useful to avoid ruptures of membranes²¹.

Chen introduced a system of entering the sinus using a turbine drill coupled with hydraulic pressure to elevate the membrane²².



Figure 1: operating RX with 2.3 mm drill close to the sinus floor.



Figure 2: Series of stops linked to the burs by a clip positioned in the crestal part of the drill.

Recently a new crestal hydraulic technique has been proposed. A particular injector, named ML, linked with a micrometric piston, Hydromab, allows for the elevation of the membrane and the filling of the sinus simultaneously^{23,24}.

The aim of this work is to propose and evaluate the efficacy of a new technique of maxillary sinus mini lift performed by aid of a turbine fitted with spherical diamond-faced burs

linked to calibrated stops that allow a reduction in the cortical thickness without entering the sinus. The cavitation produced by the simultaneous use of air and water helps the detachment of the Schneiderian membrane. The technique reduces the traumatism of operations and would decrease the percentage of failure.

Materials

Spherical diamond-faced burs are used



Figure 3: surgical approach with stop bur on the turbine and intraoperation X-ray of the bur touching the membrane.

Competing interests: None declared. Conflict of interests: None declared. All authors contributed to conception and design, manuscript preparation, read and approved the final manuscript. All authors abide by the Association for Medical Ethics (AME) ethical rules of disclosure.

(FG 201L Intensiv, Switzerland) on which calibrated stops were connected. Eight stops were used with a dimension that increased every 1 mm and allowed for a working depth of between 1 mm and 6 mm.

The stops are easy to install and change as they have a form similar to tubular rivets with a central cylindrical void of the same dimensions as the bur at the apical end and a smaller opening with 4 incisions in the metal to allow for grip on the crestal end of each stop. The bur is slid through the tubular opening on the stop and together they can be positioned on the head of the turbine (NSK Phatelus Mach-lite S, Japan) using the crestal end of the bur, as normal.

The maxillary sinus cortical grows thinner so as to obtain a hole which can enable the access to the maxillary sinus and the lifting of the Schneider membrane internally. The cavitation produced by the contemporary use of air and water helps the detachment of the schneiderian membrane.

Case study

The following case study has been described in detail to illustrate the method used. The patient was a female, 47 years old, non-smoker, healthy with no systemic disease, not on regular prescribed medication, periodontally treated, who needed an operation of maxillary sinus lift. She had no signs of inflammation of the sinus membrane. Her second upper right premolar had been extracted 4 years previous and showed approx. 4 mm residual bone height.

The operation was carried out in the following phases:

A flapless approach to the osseous crest with circular mucotome of 4.8 mm diameter (FMD Rome, Italy).

Use of a calibrated helicoidal drill with 2.3 mm diameter to approach the sinus cortical stopping one millimetre from the cortical and at the same time to recover a small quantity of

autologous bone collected in the drill coils. (Figure 1).

Preparation of spherical diamond-faced burs (FG 201L Intensiv, Switzerland) on which calibrated stops were connected (Figure 2 and Figure 3).

The membrane was detached by means of a round head compactor in order to avoid lacerations (Figure 4).




Figure 4: A compactor used to initially elevates the membrane.



Figure 5: The ML device subsequently and definitely elevates the membrane and fills the sinus with injectable biomaterial.



Figure 6: A mix of granules during the hydration.

A specific ML injector, (FMD Rome, Italy) was used to elevate the membrane and to fill the sinus^{22,23}. The biomaterial used was a mix of cortico-spongiuous porcine derived bone in an 80% collagen matrix. The granulometry is less than 300 micron (Putty by Osteobiol-Tecnoss, Italy) This was chosen due to the fact that the small

Competing interests: None declared. Conflict of interests: None declared.
All authors contributed to conception and design, manuscript preparation, read and approved the final manuscript.
All authors abide by the Association for Medical Ethics (AME) ethical rules of disclosure.

Case study

size of the granules better facilitates the injection process. (Figure 5).

A mixture was prepared and later inserted into the maxillary sinus, consisting of bone recovered by the drilling of the implant alveolus along with a mixture of cortico-spongious porcine derived bone granules with a dimension of between 250 and 1000 microns (Gen-Os by Osteobiol-Tecnoss, Italy). The graft was hydrated in a titanium dappen with saline solution (Figure 6).

A conical macro-morphology self-threading 3.5-4.8 implant (FMD, Italy; Shiner EVO with a 4.8 mm diameter crest neck) was inserted allowing for a contemporaneous horizontal expansion of the crest and a higher primary stability, thus reducing the risk of the implant accidentally sliding into the sinus. The implant apex is convex so as to avoid damaging the Schneiderian membrane (Figure 7).

Results

In the case described above ≥ 4 mm augmentations of osseous volume had been obtained. This result is comparable to those obtained in lifts where conventional techniques were applied.

Discussion

One of the most frequent reasons for failure in maxillary sinus floor lift operations is connected with the rupture of the Schneiderian membrane which, if lacerated, cannot contain the graft material. In fact the biomaterial, after becoming stable and remodeled, transforms into bone acting as fill-in and support to the implants located in the sinus.

In recent years there has been the increased tendency to use instruments and techniques so as to reduce the incidence of the type of complication mentioned above. In particular, the aim of this technique is to cut the hard tissues with extreme accuracy and minor trauma, while saving the soft tissues. The cavitation produced by the contemporary use of

air and water along with the safety aspect that the stops provide, helps the detachment of the Schneiderian membrane.

Such characteristics not only meet the requirements of maxillary sinus surgery but are also much less traumatic a more traditional technique.

In addition, there is an elimination of percussive trauma due to the exclusive use of turbine rather than the more traditional hammer technique, with a direct decrease in the patient's discomfort. Furthermore the technique reduces the number of radiographic intraoperative checks necessary during the procedure, with consequent reduction in post-operative recovery time.

Conclusion

Even with the limited number of case studies carried out to date, is it possible to foresee that the results obtained with the technique described above are encouraging.

Both the reduced percussive trauma and the low invasivity mean such a technique should be considered a valid and concrete alternative to those known and performed heretofore. Further studies are necessary in order to investigate the higher or lower efficacy in comparison with the statistically significant success.

Acknowledgment

We want to express our thanks to FMD SRL (FALAPPA MEDICAL DEVICES) for their constant and valid cooperation.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

References

1. Chiapasco M. & Romeo E. The implant-prosthesis rehabilitation in




Figure 7: The implant positioned after the sinus lift.

complex cases. Editions UTET Torino. 2003.

2. Favero GA. & Branemark P-I. Sinus lifting and osseous integration.

Stomatology and orthognatodocny Journal. Ergon 1994.

3. Summers RB. A new concept in maxillary implant surgery: The osteotomie technique. Compendium. 1994a; 15: 152-160.

4. Summers RB. The osteotome technique. Part 3- Less invasive methods of elevating the sinus floor. Compendium. 1994b; 15: 698-708.

5. Summers RB. The osteotome technique. Part 4 Future site development. Compendium. 1995; 16: 1090-09.

6. Fugazzotto PA. Sinus floor augmentation at the time of maxillary molar extraction: technique and report of preliminary results. Int J Oral Maxillofac Implants. 1999; 14: 536-542.

7. Fugazzotto PA. The modified trephine/osteotome sinus augmentation technique: Technical considerations and discussion of indications. Imp Dent. 2001; 110: 259-264.

8. Fugazzotto PA. Immediate implant placement following a modified trephine/osteotome approach: success rates of 116 implants to 4 years of function. Int.J.Oral Maxillofac. Implants 2002; 17: 113-120.

9. Fugazzotto P.A.: Sinus floor augmentation at the time of maxillary molar extraction: Success and failure rates of 137 implants in function for up to 3 years. J. Periodontol. 2002; 73: 39-44.

Case study

10. Fugazzotto P.A.: Long-term success of sinus augmentation using various surgical approaches and grafting materials. *Int.J.Oral Maxillofac. Implants* 1998; 13: 53-58.
11. Cosci F.& Luccioli M.: A new sinus lift technique in conjunction with placement of 256 implants: A 6-year retrospective study. *Imp. Dent.* 2000; 9: 363-368.
12. Malchiodi L.: *Implant surgery.* Editions Martina Bologna. 2003; 219-225.
13. Zaninari A.: Bone augmentation method using osteotomic orthodontic technique. *Dent Cadmos.* 1987 June 30;55(11):63-70.
14. Winter A.A. & Pollak A.S. & Odrich R.B.: Placement of implants in the severely atrophic posterior maxilla using localized management of the sinus floor: a preliminary study. *Int.J.Oral Maxillofac. Implants* 2002; 17: 687-695.
15. McFall T.A. , Yamane G.M., Burnett G.W.: Comparison of the cutting effect on bone of an ultrasonic cutting device and rotatory burs. *J. Oral Surg. Anesth. & Hosp. D. Serv.* 1961; 19: 200-209.
16. Horton J.E. , Tarpley T.M. Jr, Wood L.D.: Clinical applications of ultrasonic instrumentation in the surgical removal of bone. *Oral Surg. Oral Med. Oral Pathol.* 1981; 51: 236-242.
17. Vercellotti T.: Piezoelectric osseous surgery. *The modern Dentist* 2003; 5: 21-55.
18. Vercellotti T. : Piezoelectric surgery: Italian invention 2004. *Odontology Journal*, September 30, 2004.
19. Vercellotti T., De Polis S., Nevins M.: The piezoelectric bony window osteotomy and sinus membrane elevation: introduction of a new technique for simplification of the sinus augmentation procedure. *Int. J. Periodontics Restorative Dent.* 2001; 21: 561-567.
20. Vercellotti T.: Technological characteristics and clinical indications of piezoelectric bone surgery. *Minerva Stomatologica.* May 2004, volume 53, number 5.
21. E Marchetti, MA Lopez, L Confalone, S Mammolo, G Marzo.: Rialzo di seno mascellare per via crestale mediante ultrasuoni. *Journal of Osseointegration.* Ottobre 2010; 3(2)
22. Chen L, Cha J: An 8-year retrospective study: 1,100 patients receiving 1,557 implants using the minimally invasive hydraulic sinus condensing technique. *J Periodontol.* 2005 Mar;76(3):482-91
23. Andreasi Bassi, M. Lopez M.A.: J of Osteology and Biomaterials. Hydraulic sinus lift: a new method proposal. 2010; Vol. 1 N. 2: 93-101.
24. Lopez M.A., Andreasi Bassi M., Confalone L., Carinci.: Maxillary sinus floor elevation via crestal approach: the evolution of the hydraulic pressure technique, *J Craniofac Surg,* 00: 00-00, 2014.

Competing interests: None declared. Conflict of interests: None declared.
 All authors contributed to conception and design, manuscript preparation, read and approved the final manuscript.
 All authors abide by the Association for Medical Ethics (AME) ethical rules of disclosure.